

The Management of the Menopause in the light of EBM

by

Manuel Neves-e-Castro, MD

(Lisbon, Portugal)

**3rd International & 5th National Congress on
Menopause and Osteoporosis**

May 2004

Antalya / Turkey

Evidence Based Medicine

and / or

Medicine Based Evidence ?

MNC

“Without clinical expertise, practice risks becoming tyrannized by evidence.”

“Without current best evidence, practice risks becoming rapidly out of date, to the detriment of patients.”

Sackett DL et al., 1996

“Evidence alone does not make decisions”

“The new look of EBM should be **research enhanced health care**”

Haynes RB et al. Clinical expertise in the era of evidence based medicine and patient choice (Editorial) *ACP Journal Club* 2002, Mar-Apr;136(2):A11

How should we read

Epidemiology

and follow

Evidence Based Medicine ?

Information is based on

- . Observational studies
(*retrospective*)
- . Clinical trials (*prospective*)
and
- . Experimental studies
 - “in vitro”
 - “in vivo” animals
 - women

Epidemiological studies

- How were they performed?
- What similarities do they have with our clinical practice?
- How to interpret them?

The “language” of the results

- . Absolute risks
- . Relative risks
- . Number needed to treat (NNT)
- . Number needed to harm (NNH)
- . Number needed to screen (NNS)
- . Events per woman / years
- . Events per total number of women

What is

a woman /year ?!

100 woman/years = 100 women treated during 12 months

is it the same as

100 woman/years = 400 women treated during 3 months

?

Do not confuse...

Relative Risk

with

Absolute Risk!

Do not confuse...

Morbidity

with

Mortality

“Not everything that can
be counted *counts*;

and not everything that
counts can be counted”

Albert Einstein

. What **do** we know?

Researchers from the University of California, at Davis, claim **clinical trials are reported with misleading statistics**

Mayor S, *BMJ* 2002;321:1353

“Most randomised trials of new treatments published in leading medical journals (*Ann Intern Med; BMJ; JAMA; Lancet*) are reported in a potentially misleading way”.

Mayor S, *BMJ* 2002;321:1353

“Most of the trials report results based on relative risk reduction”!

“Only **18** of the papers reviewed considered absolute risk reduction”!

“Only **8** of the **359** trials reported the number needed to treat (N.N.T)”!

Mayor S, *BMJ* 2002;321:1353

Lessons from the WHI

“...most articles and broadcast segments tended to focus exclusively on either the *small absolute risks* or the *larger relative risks*, neglecting the more even-handed picture that presented both.

Since the **sharply increased *relative risks* got the most play**, news coverage about the trial's findings had an alarming cast.”

Denzer S. Editorial. *Ann Intern Med.*2003;138:352-353

“The media often failed to understand the complexities and limitations of such trials and came to its own simplified conclusions.”

“The nurse’s study and ones like it could be right and the Women’s Health Initiative could be wrong, or vice-versa”

Rossouw J, 2003

“If each is right it may be because the women in the two types of studies are different in a way that researchers have not yet figured out”.

Rossouw J, 2003

***“May be each study is wrong.
May be estrogen, in pills, is not
the chemical to focus on”***

Rossouw J, 2003

“It is quite possible that both are correct. The different results may hinge on the differences between the women who joined the studies”

Grodstein F, 2003

EBM

can be used **to do...**

and **not** **to do...**

“Annual Physical Checkup May Be an Empty Ritual”...

“Many tests that are useful, like cholesterol and blood pressure checks, need not be done every year, is said in reports to doctors, policy makers and the public”.

The New York Times; August 2003; U.S. Federal Agency for Healthcare Research and Quality..

“Annual Physical Checkup May Be an Empty Ritual “ (?!...)”

“No evidence, for example, that routine pelvic, rectal and testicular exams made any difference in overall survival rates for those with no symptoms of illness.” (!?...)

The New York Times; August 2003; U.S. Federal Agency for Healthcare Research and Quality.

CNN.com / HEALTH

SEARCH

The Web  CNN.com 

Search

Home Page

World

U.S.

Weather

Business

Sports

Politics

Law

Technology

Science & Space

Health

Entertainment

Travel

Education

Special Reports



SERVICES

Video

E-Mail Services


CNN To Go

Study: Fellatio may significantly decrease the risk of breast cancer in women

Thursday, October 2, 2003 Posted: 9:19 AM EDT (1319 GMT)

(AP) -- Women who perform the act of fellatio on a regular basis, one to two times a week, may reduce their risk of breast cancer by up to 40 percent, a North Carolina State University study found.

Doctors had never suspected a link between the act of fellatio and breast cancer, but new research being performed at North Carolina State University is starting to suggest that there could be an important link between the two.



Story Tools

  SAVE THIS  EMAIL THIS

Study: *”Fellatio may significantly decrease the risk of breast cancer in women” !...*

(AP) – “Women who perform the act of fellatio on a regular basis, one to two times a week, may reduce their risk of breast cancer by up to 40 percent, a North Carolina State University study found”

Thursday, October 2, 2003 Posted: 9:19 AM EDT (1319 GMT)

“CNN.com./HEALTH”

therefore ...

***When this was read and heard
all over the World...***

**Most likely, some people
started thinking that the
seminal plasma might contain
an *orally active* protective
molecule ...**

however...

It was soon found out that **it had been a joke** sent to the Web by one **ironical** N.C. University student ,who wanted to have fun with some epidemiologists! ...

The Truth?

*“The objective of both basic and clinical science is **to know the truth**”.*

*“Every epidemiologic study, no matter how good or how large, **gives only one view of the truth**”.*

“It takes many views to come close to seeing the truth”

Bush TL. *Int J Fertil.*2001;46:56-59